



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

load. It is constructed almost entirely of heavy gaspipe, and is extremely rigid. The polar axis is hollow, and is fitted with an eyepiece, as an aid in adjusting. The automatic arrangements for securing certainty in exposing have been elaborated upon to such an extent that the inventor now feels certain that nothing can go wrong, at least with this part of the expedition.

Professor BURCKHALTER'S parting injunction to his friends was not to wish him a good time or a pleasant journey, but that he might have two minutes of clear sky at the right time. Those of us who have heard him describe the disappointing day in Japan in 1896, and who realize what the success of this expedition means, will be certain to remember him on the eventful day. If friendly good wishes can insure success, he will have it.

ALLEN H. BABCOCK.

ELEMENTS OF COMET *b*, 1897 (PERRINE).

From Mt. Hamilton observations, made on October 16th, 18th, and 20th, we have computed the following elements of the orbit of this comet:—

$$\begin{aligned} T &= 1897 \text{ Dec. } 9.89171 \text{ G. M. T.} \\ \omega &= 67^\circ 6' 55''.2 \\ \Omega &= 32^\circ 8' 37'' .4 \\ i &= 69^\circ 45' 43'' .2 \end{aligned} \left. \begin{array}{l} \text{Mean equinox and ecliptic} \\ \text{of } 1897.0 \end{array} \right\}$$

$\log q = 0.129500$.

Residuals for the middle place (O—C):—

$$\Delta\lambda \cos \beta = +2''.8, \Delta\beta = +2''.4.$$

A comparison of observations made on November 1st with the ephemeris positions computed from these elements shows a satisfactory agreement. W. J. HUSSEY and R. G. AITKEN.

November 3, 1897.

ASTRONOMICAL TELEGRAMS (*Translations*).

Lick Observatory, Oct. 17, 1897.

To Harvard College Observatory: } (Sent 1:00 A. M.)
To Students' Observatory, Berkeley: }

A comet was discovered by C. D. PERRINE, October 16.7398, G. M. T.; R. A. $3^h 36^m 7\overset{.}{s} .6$; N. P. D. $23^\circ 13' 16''$. The comet is about $2'$ in diameter, is as bright as an eighth magnitude star, has a well-defined nucleus and a tail less than $30'$ long.

Lick Observatory, Oct. 18, 1897.

To Harvard College Observatory: } (Sent 1:40 P.M.)
To Students' Observatory, Berkeley: }

Comet *b*, 1897 (PERRINE), was observed by C. D. PERRINE,
October 17.7121, G. M. T.; R. A. $3^h 30^m 25^s.7$; N. P. D. 21°
 $42' 47''$.

Lick Observatory, Oct. 18, 1897.

To Harvard College Observatory: } (Sent 9:55 P.M.)
To Students' Observatory, Berkeley: }

Comet *b*, 1897 (PERRINE), was observed by W. J. HUSSEY,
October 18.6498, G. M. T.; R. A. $3^h 24^m 2^s.2$; N. P. D. 20°
 $16' 06''$.

Lick Observatory, Oct. 19, 1897.

To Harvard College Observatory: (Sent 5:10 P.M.)

Comet *b*, 1897, was observed with the Meridian Circle by
R. H. TUCKER, October 18.9011, G. M. T.; R. A. $3^h 22^m 5^s.5$;
N. P. D. $19^\circ 52' 49''$.

Lick Observatory, Oct. 19, 1897.

To Harvard College Observatory: (Sent 9:15 A.M.)

Elements and ephemeris of Comet *b*, 1897, were computed by
W. J. HUSSEY and R. G. AITKEN.

ELEMENTS.

$T = G. M. T. 1897, Dec. 9.2300.$

$\omega = 66^\circ 28'$ }
 $\Omega = 32^\circ 5'$ }
 $i = 69^\circ 38'$ } Mean equinox of 1897.0
 $q = 1.3525.$

[The ephemeris at four-day intervals, from October 20th to
November 1, 1897, is here omitted.]